

SHEP 92/93-11

hep-ph/9302202

Supersymmetric Higgs Bosons at the Limit

T. Elliott, S. F. King and P. L. White,

Physics Department,

University of Southampton,

Southampton,

SO9 5NH,

U.K.

Abstract

We obtain an improved upper bound on the lightest neutral CP-even Higgs boson mass from a low energy renormalisation-group analysis of the Higgs sector of the next-to-minimal supersymmetric standard model. We find $m_h < 145$ GeV for $m_t = 90$ GeV, decreasing to $m_h < 123$ GeV for $m_t = 180$ GeV. We also discuss the light Higgs spectrum in the region of parameter space close to the upper bound.

The question of the origin of electroweak mass is one of the most urgent questions of present day particle physics. The discovery of a particle which resembles the Higgs boson of the minimal standard model, and the measurement of its mass, will provide clues as to the nature of new physics beyond the standard model. In this letter we shall be concerned with the question of how heavy the lightest neutral CP-even supersymmetric Higgs boson, h^0 , can be within the framework of supersymmetric grand unified theories (SUSY GUTs) [1]. In SUSY GUTs all the Yukawa couplings are constrained to remain perturbative in the region $M_{SUSY} \sim 1$ TeV to $M_{GUT} \sim 10^{16}$ GeV. This constraint provides a maximum value at low energies for those Yukawa couplings which are not asymptotically-free, and is obtained from the renormalisation group (RG) equations together with the boundary conditions that the couplings become non-perturbative at M_{GUT} – the so-called “triviality limit”. In the minimal supersymmetric standard model (MSSM) [2], the triviality limits provide a useful bound on the top quark mass m_t . The upper bound on the h^0 mass, m_h , in the MSSM, including radiative corrections, has recently been the subject of much discussion[3, 4, 5, 6, 7, 8, 9]. However the MSSM is not the most general low energy manifestation of SUSY GUTs. It is possible that SUSY GUTs give rise to a low energy theory which contains an additional gauge singlet field, the so called next-to-minimal supersymmetric standard model (NMSSM) [10, 11, 12]. Here we shall concentrate on the question of the upper-bound on m_h in the NMSSM which is obtained from triviality limits of Yukawa couplings.

In the NMSSM there are two Higgs doublets H_1, H_2 and a complex Higgs singlet N , leading to three neutral CP-even Higgs bosons, two neutral CP-odd states and two charged bosons in the physical spectrum. The superpotential has the form

$$W = h_t Q t^c H_2 + \lambda N H_1 H_2 - \frac{k}{3} N^3 + \dots, \quad (1)$$

where the superfield Q contains the left-handed third family quark doublet (t_L, b_L) ,

the superfield t^c contains the charge conjugate of the right-handed top quark field, $H_{1,2}$ contain the Higgs doublets and N contains the Higgs gauge singlet. The ellipsis represents terms whose relatively small couplings will not play a role in our analysis. The NMSSM removes the $\mu H_1 H_2$ term in the superpotential of the MSSM and replaces its effect by the vacuum expectation value (VEV) of the Higgs singlet, $\langle N \rangle = x$. The other VEVs are $\langle H_{1,2} \rangle = v_{1,2}$, where $v = \sqrt{v_1^2 + v_2^2} = 174$ GeV. The trilinear term $\frac{k}{3}N^3$ is necessary in order to avoid an approximate global $U(1)$ symmetry (broken by instanton effects) which, when spontaneously broken by the VEVs, would lead to an axion. The Higgs sector of the NMSSM reduces to that of the MSSM in the limit $r = x/v \rightarrow \infty$ with λx and kx fixed, where one of the neutral CP-even states and one of the CP-odd states decouple [11].

An upper bound on the lightest neutral CP-even scalar h^0 in the NMSSM may be obtained from the real symmetric 3×3 neutral scalar mass squared matrix, by using the fact that the smallest eigenvalue of such a matrix must be smaller than the smallest eigenvalue of its upper 2×2 block. The resulting bound at tree-level is [12]

$$m_h^2 \leq M_Z^2 + (\lambda^2 v^2 - M_Z^2) \sin^2 2\beta. \quad (2)$$

where $\tan \beta \equiv \frac{v_2}{v_1}$, and λ is regarded as a running parameter evaluated at M_{SUSY} . The upper bound on m_h is determined by the maximum value of $\lambda(M_{SUSY})$, henceforth denoted λ_{max} . The value of λ_{max} is obtained by solving the SUSY RG equations for the Yukawa couplings h_t , λ and k in the region $M_{SUSY} = 1$ TeV to $M_{GUT} = 10^{16}$ GeV [13, 14]. If the Yukawa couplings h_t , λ and k are all initially large at M_{GUT} then they approach low energy fixed point ratios [13]. However, if the boundary condition at M_{GUT} is $\lambda \gg k$, then larger values of $\lambda(M_{SUSY})$ can be achieved. We have repeated the calculation of ref.[14] and found that for $h_t(M_{SUSY}) = 0.5-1.0$, $\lambda_{max} = 0.87-0.70$ and for $h_t(M_{SUSY}) \rightarrow 1.06$, $\lambda_{max} \rightarrow 0$ (with $k = 0$ always). Radiative corrections to

¹ $h_t(M_{SUSY}) \leq 1.06$ is the triviality bound, which, together with $m_t = h_t(m_t)v \sin \beta$, where $h_t(m_t) \leq 1.12$, implies the bound $m_t \leq 195$ GeV.

the tree-level bound in Eq.(2) have been considered in refs.[15, 16, 17]. In ref.[16] these were estimated from a low energy RG analysis of the Higgs sector of the model between M_{SUSY} and a lower scale μ , assuming that only one Higgs boson has a mass below M_{SUSY} . Here we shall consider the more general case in which both Higgs doublets and the Higgs singlet may be lighter than M_{SUSY} , making the usual approximation of hard decoupling below M_{SUSY} of the superpartners. We shall then apply this technique to obtain a new upper bound on m_h , and supplement the discussion with a full numerical analysis of the light Higgs spectrum of the NMSSM in the region of parameter space close to this upper bound.

At energy scales above M_{SUSY} the Higgs potential in the NMSSM with general soft SUSY breaking terms is given from Eq.(1) by

$$\begin{aligned}
V_{Higgs} = & \lambda^2 \left[(|H_1|^2 + |H_2|^2)|N|^2 + |H_1 H_2|^2 \right] + k^2 |N|^4 - \lambda k (H_1 H_2 N^{*2} + H.c.) \\
& + \frac{1}{8} (g_1^2 + g_2^2) (|H_2|^2 - |H_1|^2)^2 + \frac{1}{2} g_2^2 |H_2^\dagger H_1|^2 \\
& + m_{H_1}^2 |H_1|^2 + m_{H_2}^2 |H_2|^2 + m_N^2 |N|^2 \\
& - \lambda A_\lambda (H_1 H_2 N + H.c.) - \frac{1}{3} k A_k (N^3 + H.c.), \tag{3}
\end{aligned}$$

where $H_1 H_2 = H_1^T i \sigma_2 H_2$, σ_2 is the second Pauli matrix and $H_1^T = (H_1^0 \ H_1^-)$, $H_2^T = (H_2^+ \ H_2^0)$. The parameters m_{H_i} , m_N , $A_{\lambda,k}$ are associated with soft SUSY breaking terms.² We shall take the parameters λ , k , A_λ and A_k to be real and positive, which is a sufficient condition for the vacuum to conserve CP and leads to a choice of vacuum in which all the VEVs x, v_1, v_2 are real and positive [11]. Note that the range of the parameters is restricted by the condition that the vacuum does not break QED, which is not automatic in the NMSSM, and is equivalent to the condition that $m_c^2 \geq 0$, where m_c is the mass of the physical charged Higgs $H^\pm$. Also it is not automatic that the vacuum does not break QCD although it has been checked that this is the case for values of r and $\tan \beta$ in the range 0.05-20 [11]. Finally, the range of

²In our analysis we shall not consider any other soft SUSY breaking parameters.

the parameter A_k is restricted by the requirements that $\langle V_{Higgs} \rangle < 0$, and all mass squared eigenvalues are positive. We shall use the above conditions in our analysis.

The effective theory below M_{SUSY} is just the standard model with two light Higgs doublets and a light Higgs singlet. Thus the Higgs potential at some low energy scale $\mu < M_{SUSY}$ is given by the general expression ³

$$\begin{aligned}
V_{Higgs} = & \frac{1}{2}\lambda_1(H_1^\dagger H_1)^2 + \frac{1}{2}\lambda_2(H_2^\dagger H_2)^2 + (\lambda_3 + \lambda_4)(H_1^\dagger H_1)(H_2^\dagger H_2) \\
& - \lambda_4|H_2^\dagger H_1|^2 + \lambda_5|N|^2|H_1|^2 + \lambda_6|N|^2|H_2|^2 \\
& + \lambda_7(N^{*2}H_1H_2 + H.c.) + \lambda_8|N|^4 \\
& + m_1^2|H_1|^2 + m_2^2|H_2|^2 + m_3^2|N|^2 \\
& - m_4(H_1H_2N + H.c.) - \frac{1}{3}m_5(N^3 + H.c.).
\end{aligned} \tag{4}$$

Comparing Eqs.(3) and (4), the running quartic couplings λ_i and the mass parameters m_i must satisfy the following boundary conditions at M_{SUSY}

$$\begin{aligned}
\lambda_1 &= \lambda_2 = \frac{1}{4}(g_2^2 + g_1^2), \quad \lambda_3 = \frac{1}{4}(g_2^2 - g_1^2) \\
\lambda_4 &= \lambda^2 - \frac{1}{2}g_2^2, \quad \lambda_5 = \lambda_6 = \lambda^2, \quad \lambda_7 = -\lambda k, \quad \lambda_8 = k^2, \\
m_1 &= m_{H_1}, \quad m_2 = m_{H_2}, \quad m_3 = m_N, \\
m_4 &= \lambda A_\lambda, \quad m_5 = k A_k.
\end{aligned} \tag{5}$$

At energy scales μ below M_{SUSY} , the values of the quartic couplings may be obtained by solving the following RG equations which we have derived for the potential in Eq.(4) with the aid of the general results in ref.[18]:

$$\begin{aligned}
16\pi^2 \frac{\partial \lambda_1}{\partial t} &= 12\lambda_1^2 + 4\lambda_3^2 + 4\lambda_3\lambda_4 + 2\lambda_4^2 + 2\lambda_5^2 \\
&- \lambda_1(3g_1^2 + 9g_2^2) + \frac{3}{4}g_1^4 + \frac{9}{4}g_2^4 + \frac{3}{2}g_1^2g_2^2, \\
16\pi^2 \frac{\partial \lambda_2}{\partial t} &= 12\lambda_2^2 + 4\lambda_3^2 + 4\lambda_3\lambda_4 + 2\lambda_4^2 + 2\lambda_6^2
\end{aligned}$$

³There is a global $U(1)$ symmetry of the quartic potential which forbids the radiative generation of other terms not included in Eq.(4).

$$\begin{aligned}
& - \lambda_2(3g_1^2 + 9g_2^2) + \frac{3}{4}g_1^4 + \frac{9}{4}g_2^4 + \frac{3}{2}g_1^2g_2^2 \\
& + 12h_t^2\lambda_2 - 12h_t^4, \\
16\pi^2\frac{\partial\lambda_3}{\partial t} &= 2(\lambda_1 + \lambda_2)(3\lambda_3 + \lambda_4) + 4\lambda_3^2 + 2\lambda_4^2 - 2\lambda_5\lambda_6 \\
& - \lambda_3(3g_1^2 + 9g_2^2) + \frac{3}{4}g_1^4 + \frac{9}{4}g_2^4 - \frac{3}{2}g_1^2g_2^2 + 6h_t^2\lambda_3, \\
16\pi^2\frac{\partial\lambda_4}{\partial t} &= 2\lambda_4(\lambda_1 + \lambda_2 + 4\lambda_3 + 2\lambda_4) + 4\lambda_7^2 \\
& - \lambda_4(3g_1^2 + 9g_2^2) + 3g_1^2g_2^2 + 6h_t^2\lambda_4, \\
16\pi^2\frac{\partial\lambda_5}{\partial t} &= 2\lambda_5(3\lambda_1 + 2\lambda_5 + 4\lambda_8) + 2\lambda_6(2\lambda_3 + \lambda_4) + 8\lambda_7^2 \\
& - \frac{1}{2}\lambda_5(3g_1^2 + 9g_2^2), \\
16\pi^2\frac{\partial\lambda_6}{\partial t} &= 2\lambda_5(2\lambda_3 + \lambda_4) + 2\lambda_6(3\lambda_2 + 2\lambda_6 + 4\lambda_8) + 8\lambda_7^2 \\
& - \frac{1}{2}\lambda_6(3g_1^2 + 9g_2^2) + 6h_t^2\lambda_6, \\
16\pi^2\frac{\partial\lambda_7}{\partial t} &= 2\lambda_7(\lambda_3 + 2\lambda_4 + 2\lambda_5 + 2\lambda_6 + 2\lambda_8) \\
& - \frac{1}{2}\lambda_7(3g_1^2 + 9g_2^2) + 3h_t^2\lambda_7, \\
16\pi^2\frac{\partial\lambda_8}{\partial t} &= 2\lambda_5^2 + 2\lambda_6^2 + 4\lambda_7^2 + 20\lambda_8^2, \tag{6}
\end{aligned}$$

where $t = \ln \mu$, and h_t obeys the RG equation

$$16\pi^2\frac{\partial h_t}{\partial t} = \frac{9}{2}h_t^3 - h_t(8g_3^2 + \frac{9}{4}g_2^2 + \frac{17}{12}g_1^2). \tag{7}$$

The gauge couplings g_i obey the RG equations of the standard model with two Higgs doublets and three fermion families below M_{SUSY} , namely

$$16\pi^2\frac{\partial g_i}{\partial t} = -c_i g_i^3 \quad (c_1 = -7, c_2 = 3, c_3 = 7), \tag{8}$$

and we shall take $\alpha_{1,2,3}(M_Z) = 0.0102, 0.0336, 0.113$, respectively.

The minimisation conditions implied by $\frac{\partial V_{Higgs}}{\partial v_i} = 0$ and $\frac{\partial V_{Higgs}}{\partial x} = 0$ allow us to eliminate the low energy parameters m_1, m_2, m_3 . The remaining masses m_4 and m_5 are related to the parameters A_λ and A_k at M_{SUSY} by Eq.(5). Below this scale we shall regard m_4 and m_5 as free parameters. The charged Higgs squared mass is given

by

$$m_c^2 = \frac{2x}{\sin 2\beta}(m_4 - \lambda_7 x) - \lambda_4 v^2, \quad (9)$$

which allows us to eliminate the parameter m_4 in favour of the more physical parameter m_c^2 . The physical neutral CP-odd pseudoscalar mass squared symmetric matrix elements are

$$\begin{aligned} (M_{ps}^2)_{11} &= 3xm_5 + \frac{v^2}{4x^2} \sin^2 2\beta (m_c^2 + \lambda_4 v^2) - \frac{3}{2} \lambda_7 v^2 \sin 2\beta, \\ (M_{ps}^2)_{12} &= \frac{v}{2x} \sin 2\beta (m_c^2 + \lambda_4 v^2) + 3vx\lambda_7, \\ (M_{ps}^2)_{22} &= m_c^2 + \lambda_4 v^2. \end{aligned} \quad (10)$$

The neutral CP-even scalar mass squared symmetric matrix elements are, in the basis $1, 2, 3 = H_1, H_2, N$,

$$\begin{aligned} (M_s^2)_{11} &= 2\lambda_1 v_1^2 + (m_c^2 + \lambda_4 v^2) \sin^2 \beta, \\ (M_s^2)_{12} &= \sin 2\beta [(\lambda_3 + \frac{1}{2}\lambda_4)v^2 - \frac{1}{2}m_c^2], \\ (M_s^2)_{22} &= 2\lambda_2 v_2^2 + \cos^2 \beta (m_c^2 + \lambda_4 v^2), \\ (M_s^2)_{13} &= x(2\lambda_5 v_1 + \lambda_7 v_2) - \frac{v_2}{2x} \sin 2\beta (m_c^2 + \lambda_4 v^2), \\ (M_s^2)_{23} &= x(2\lambda_6 v_2 + \lambda_7 v_1) - \frac{v_1}{2x} \sin 2\beta (m_c^2 + \lambda_4 v^2), \\ (M_s^2)_{33} &= 4\lambda_8 x^2 + \frac{v^2}{4x^2} \sin^2 2\beta (m_c^2 + \lambda_4 v^2) + \frac{1}{2} \lambda_7 v^2 \sin 2\beta - xm_5. \end{aligned} \quad (11)$$

The model is then specified by the 6 parameters $\lambda, k, \tan \beta = v_2/v_1, r = x/v, m_5$, and m_c^2 . Our detailed procedure for evaluating Higgs masses is as follows. Given the gauge couplings $g_i(M_Z)$, we find $g_i(m_t)$ by solving the RG equations (8). For a given m_t and $\tan \beta$ we obtain $h_t(m_t)$, then use its RG equation (7) and those for the gauge couplings (8) to evaluate $h_t(M_{SUSY})$ and $g_i(M_{SUSY})$. Given these couplings and some choice of λ and k we then use the boundary conditions in Eq.(5) and the RG equations (6),(7),(8), to find $\lambda_i(\mu)$, where μ is close to mass-shell for the Higgs

boson of interest.⁴ In practice we shall take $M_{SUSY} = 1$ TeV, $\mu = 150$ GeV. The neutral Higgs masses are then calculated from Eqs.(10),(11) with $\lambda_i = \lambda_i(\mu)$. The diagonalisation of these matrices also gives the mixing angles, so that the amplitude of the singlet state N in each of the neutral mass eigenstates may be calculated.

We shall now obtain an upper bound on the mass of the lightest neutral scalar Higgs boson h^0 , by using the fact that m_h^2 must not exceed the lower eigenvalue of the upper 2×2 block matrix, $(M_s^2)_{ij}$, $i, j = 1, 2$. The resulting upper bound is a complicated function of m_c^2

$$m_h^2 \leq \frac{1}{2}(A + m_c^2) - \frac{1}{2}\sqrt{(m_c^2 + B)^2 + C^2 - B^2}, \quad (12)$$

where

$$\begin{aligned} A &= v^2(\lambda_1 + \lambda_2 + \lambda_4) + v^2(\lambda_1 - \lambda_2) \cos 2\beta, \\ B &= -v^2 \cos 2\beta[(\lambda_1 - \lambda_2) + (\lambda_1 + \lambda_2 - \lambda_4) \cos 2\beta] - 2v^2 \sin^2 2\beta(\lambda_3 + \frac{1}{2}\lambda_4), \\ C^2 &= v^4[(\lambda_1 - \lambda_2) + (\lambda_1 + \lambda_2 - \lambda_4) \cos 2\beta]^2 + 4v^4 \sin^2 2\beta(\lambda_3 + \frac{1}{2}\lambda_4)^2. \end{aligned} \quad (13)$$

It is easy to show that the above bound reaches a maximum asymptotically for $m_c \rightarrow \infty$. Since $(C^2 - B^2) \geq 0$, we obtain the m_c -independent bound

$$m_h^2 \leq \frac{1}{2}(A - B). \quad (14)$$

Inserting A, B from Eq.(13) and using the boundary conditions in Eq.(5) leads to an upper bound of the form

$$\begin{aligned} m_h^2 &\leq M_Z^2 + (\lambda^2 v^2 - M_Z^2) \sin^2 2\beta \\ &+ \frac{v^2}{2}[(\delta\lambda_1 + \delta\lambda_2 + 2\delta\lambda_3 + 2\delta\lambda_4) + 2(\delta\lambda_1 - \delta\lambda_2) \cos 2\beta \\ &+ (\delta\lambda_1 + \delta\lambda_2 - 2\delta\lambda_3 - 2\delta\lambda_4) \cos^2 2\beta], \end{aligned} \quad (15)$$

⁴In integrating the RG equations from M_{SUSY} down to μ we have decoupled the terms involving h_t below the scale m_t , but we have not decoupled any Higgs bosons below their mass scales. We would not expect such a decoupling to change our results significantly providing there are no very heavy scalars.

where $\delta\lambda_i = \lambda_i(\mu) - \lambda_i(M_{SUSY})$, and, strictly speaking, M_Z should be evaluated at M_{SUSY} . Using the RG equations (6), keeping only terms quadrilinear in λ , h_t , and k and making a small $\delta\lambda_i$ approximation, Eq.(15) yields the analytic bound ⁵

$$\begin{aligned}
m_h^2 &\leq M_Z^2 + (\lambda^2 v^2 - M_Z^2) \sin^2 2\beta \\
&+ \frac{v^2}{32\pi^2} \ln \left(\frac{M_{SUSY}}{\mu} \right) [(12h_t^4 - 12\lambda^2 h_t^2 - 8\lambda^2 k^2 - 24\lambda^4) - 24h_t^4 \cos 2\beta \\
&+ (12h_t^4 + 12\lambda^2 h_t^2 + 8\lambda^2 k^2 + 8\lambda^4) \cos^2 2\beta].
\end{aligned} \tag{16}$$

In Eq.(16), the tree level bound in Eq.(2) is corrected by a function of h_t , $\cos 2\beta$, λ and k . It is easy to show that this function is maximised for $\lambda > \lambda_{max}$ and $k = 0$, and so we shall use λ_{max} and $k = 0$, as in the case of the tree level bound discussed previously. With this information in hand, let us now return to the more exact result in Eq.(15) in which the values of $\delta\lambda_i$ are obtained from numerically integrating the RG equations, decoupling the top quark below its mass and choosing $\mu = 150$ GeV. For each value of m_t , we have determined the value of $h_t(M_{SUSY})$ and the corresponding value of λ_{max} which maximise the r.h.s. of Eq.(15) from a numerical analysis of the triviality condition as discussed previously. These are shown in Fig.1, together with $\sin \beta$ which is eliminated from Eq.(15) using $m_t = h_t(m_t)v \sin \beta$.

Using the parameters in Fig.1, we obtain the upper bound on the lightest neutral Higgs scalar in the NMSSM as a function of m_t as shown in Fig.2. We again emphasise that we have considered the more general case of two light Higgs doublets plus a light singlet all below the scale M_{SUSY} , whereas previous results [16, 17] have only considered the case of one light Higgs boson. From Fig.2 we find that for $m_t = 90$ GeV, $m_h \leq 145$ GeV, which agrees to within 2 GeV with both refs.[16, 17]. Our bound for larger top quark masses, although qualitatively similar to refs.[16, 17], can differ by several GeV. It can easily be checked that the analytic formula in Eq.(16) agrees with the full numerical bound shown to within 5 GeV across the entire m_t

⁵Note that Eq.(16) reduces to the bound in the MSSM in the limit that $\lambda \rightarrow 0$.

region, giving a bound somewhat lower than the true bound for $m_t = 90 - 160$ GeV, somewhat higher for $m_t = 180 - 190$ GeV, and in good agreement for $m_t \approx 170$ GeV. If the λ dependent terms in the radiative corrections in Eq.(16) were dropped, this would produce a bound about 7 GeV higher than the true bound over the entire m_t range.

We shall now discuss the physical parameter regions in which the upper bound on m_h in Fig.2 is approximately realised, and discuss the light Higgs boson spectrum in these regions. As mentioned earlier, the m_c independent bound in Eqs.(15),(16) is exactly valid only asymptotically, but actually the bound rapidly approaches its maximum for large values of m_c of the order of the bound itself. Since it is known that the allowed range of m_c is controlled by r and that larger values of m_c are associated with the larger values of r , we shall restrict ourselves to $r = 1.0$ and 10.0 , as shown in Figs.3 and 4.

Fig.3(a) shows the light Higgs boson spectrum as a function of m_c for $m_t = 150$ GeV, for $r = 1.0$, where $\tan \beta = 1.7$ is from Fig. 1. We have taken $\lambda = 0.65$ and $k = 0.1$, rather than $\lambda_{max} = 0.7$ and $k = 0$ since these values would imply a massless CP-odd scalar, which is phenomenologically ruled out. We have also taken $m_5 = 0$. Only the lightest neutral CP-even scalars (solid) and the lightest CP-odd pseudoscalar (dashed) are included in Fig.3(a). The lightest scalar mass m_h reaches a maximum of $m_h = 88$ GeV. In Fig.3(b) we show the N components of the bosons in Fig.3(a). This is of interest since the N component is a gauge singlet, and thus decoupled from the gauge bosons. The lightest CP-even scalar is about 80% decoupled rising to being almost 100% decoupled at its maximum. It is obvious that the bound which follows from the upper 2×2 block matrix can only approach saturation if the full scalar matrix is approximately block diagonal. Since the lightest scalar is approximately decoupled near its maximum mass, the physically observable second lightest scalar must respect the bound at this point. Taking m_5 to be non-zero and positive (negative) serves to

lower (raise) $(M_s^2)_{33}$ (identified as the decoupled scalar mass) while leaving the other elements unchanged, as is clear from Eq.(11). Finally it is clear from Figs.3(a),3(b) that the lightest CP-odd state presents no phenomenological problems, since as well as being heavy it is over 90% decoupled over the entire m_c range.

In Fig.4(a) we show the light Higgs spectrum for $r = 10.0$, choosing the other parameters to be the same as in Fig.3(a). The relevant range of m_c is now 2.3-2.5 TeV which is strictly outside the range of validity of our calculation, since we do not consider Higgs decoupling and choose $\mu = 150$ GeV (of order m_h) as in Fig.3(a). Nevertheless Fig.4(a) shows that the lightest CP-even scalar mass reaches a maximum of 118 GeV, close to the minimum of the physically observable second lightest scalar in Fig.3(a). Fig.4(b) shows that in this case the lightest scalar is the detectable one, being almost 0% decoupled at its maximum, while the second lightest scalar is almost 100% decoupled at this point. The CP-odd boson in Fig.4 is now nearly 100% decoupled over the whole range of m_c .

In conclusion, the lightest CP-even neutral Higgs boson in the NMSSM must respect the bounds, $m_h < 145$ GeV for $m_t = 90$ GeV, decreasing to $m_h < 123$ GeV for $m_t = 180$ GeV. Our analysis is based on the assumption of a SUSY desert between $M_{SUSY} = 1$ TeV and $M_{GUT} = 10^{16}$ GeV, and involves the assumption of hard decoupling of the superpartners. However, our calculations do include the effects of other light Higgs bosons which previous analyses have ignored. We have presented an analytic expression in Eq.(16) for the radiative corrections in the NMSSM, which agrees well with our full numerical bound in Fig.2. In addition we have discussed the light Higgs boson spectrum in a region of parameter space which approximately realises the bound, and have seen that there are indeed other light Higgs bosons whose effects must be taken into account. For $r \sim 1$ (Fig.3), the lightest CP-even scalar is not very strongly physically coupled when it is close to its maximum mass, and the second lightest CP-even scalar, which must also respect the bound, may be easier to

discover.

Acknowledgements

TE and SFK would like to thank the SERC for financial support.

References

- [1] H. P. Nilles, Phys. Rep. 110 (1984) 1; H. E. Haber and G. L. Kane, Phys. Rep. 117 (1985) 75.
- [2] J. F. Gunion, H. E. Haber, G. L. Kane and S. Dawson, “The Higgs Hunter’s Guide” (Addison-Wesley, Reading, MA, 1990)
- [3] J. Ellis, G. Ridolfi and F. Zwirner, Phys. Lett. B257 (1991) 83; *ibid* B262 (1991) 477; A. Brignole, J. Ellis, G. Ridolfi and F. Zwirner, Phys. Lett. B271 (1991) 123.
- [4] H. Haber and R. Hempfling, Phys. Rev. Lett.66 (1991) 1815.
- [5] Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. 85 (1991)1; *ibid* Phys. Lett. B262 (1991) 54.
- [6] R. Barbieri, M. Frigeni and F. Caravaglios, Phys. Lett. B258 (1991) 167.
- [7] J. L. Lopez and D. V. Nanopoulos, Phys. Lett. B266 (1991) 397.
- [8] A. Brignole, Phys. Lett. B277 (1992) 313; *ibid* Phys. Lett. B281 (1992) 284.
- [9] K. Sasaki, M. Carena and C. E. M. Wagner, Nucl. Phys. B381 (1992) 66.
- [10] P. Fayet, Nucl. Phys. B90 (1975) 104.
- [11] J. Ellis, J. F. Gunion, H. E. Haber, L. Roszkowski and F. Zwirner, Phys. Rev. D39 (1989) 844.
- [12] L. Durand and J. L. Lopez, Phys. Lett. B217 (1989) 463; M. Drees, Int. J. Mod. Phys. A4 (1989) 3635.
- [13] P. Binetruy and C. A. Savoy, Phys. Lett. B277 (1992) 453.

- [14] J. R. Espinosa and M Quiros, Phys. Lett. B279 (1992) 92.
- [15] U. Ellwanger and M. Rausch de Traubenberg, Z. Phys. C53 (1992) 521; U. Ellwanger and M. Lindner, Heidelberg preprint HD-THEP-92-48.
- [16] W. ter Veldhuis, Purdue preprint PURD-TH-92-11.
- [17] J. R. Espinosa and M. Quiros, Madrid preprint IEM-FT-64/92 and hep-ph/9212305.
- [18] T. P. Cheng, E. Eichten and Ling-Fong Li, Phys. Rev. D9 (1974) 2259.

Figure Captions

Figure 1 : The values of h_t (solid), λ_{max} (short dashed) and $\sin \beta$ (long dashed) realising the upper bound on m_h in Figure 2.

Figure 2 : The full numerical bound on the lightest neutral CP-even Higgs scalar as a function of m_t .

Figure 3(a) : The light Higgs boson spectrum for the parameters $m_t=150$ GeV, $r = 1.0$, $\tan \beta = 1.7$, $\lambda = 0.65$, $k = 0.1$. The solid lines are CP-even scalar masses, and the dashed line is the CP-odd mass.

Figure 3(b) : The N components of each of the bosons in Fig.3(a) where the solid line is for the lightest CP-even scalar, the dashed line is for the second lightest CP-even scalar, and the dotted line is for the CP-odd pseudoscalar.

Figure 4(a) : The light Higgs boson spectrum for $r = 10.0$. The other parameters are as in Figure 3(a). The solid lines are CP-even scalar masses, and the dashed line is the CP-odd mass.

Figure 4(b) : The N components of each of the bosons in Fig.4(a) where, as in Fig.3(b), the solid line is for the lightest CP-even scalar, the dashed line is for the second lightest CP-even scalar, and the dotted line is for the CP-odd pseudoscalar.

Figure 1




Figure 2




Figure 3.a




Figure 3.b




Figure 4.a




Figure 4.b

